

TITLE

Women's dietary changes before and during pregnancy: A systematic review

AUTHOR

Hillier, Sarah E. and Olander, Ellinor K.

JOURNAL

Midwifery

DATE DEPOSITED

14 June 2017

This version available at

<https://research.stmarys.ac.uk/id/eprint/1532/>

COPYRIGHT AND REUSE

Open Research Archive makes this work available, in accordance with publisher policies, for research purposes.

VERSIONS

The version presented here may differ from the published version. For citation purposes, please consult the published version for pagination, volume/issue and date of publication.

Title: Women's dietary changes before and during pregnancy: a systematic review

Authors: Sarah E. Hillier^{1,3} and Ellinor K. Olander²

Affiliations: ¹School of Health and Applied Sciences, St Mary's University, Twickenham, London

²Centre for Maternal and Child Health Research, City University, London

³Department of Sport and Health Sciences, Faculty of Health and Life Sciences,
Oxford Brookes University, Oxford

Corresponding Author: Dr Sarah E Hillier

Lecturer in Nutrition

Department of Sport and Health Sciences

Faculty of Health and Life Sciences

Oxford Brookes University

Oxford

sarahhillier@brookes.ac.uk

Abstract**Background**

Dietary intake before and during pregnancy has significant health outcomes for both mother and child, including a healthy gestational weight gain. To ensure effective interventions are successfully developed to improve dietary intake during pregnancy, it is important to understand what dietary changes pregnant women make without intervention.

Aims

To systematically identify and review studies examining women's dietary changes before and during pregnancy and to identify characteristics of the women making these changes.

Methods

A systematic search strategy was employed using three databases (Web of Science, CINAHL and PubMed) in May 2016. Search terms included those relating to preconception, pregnancy and diet. All papers were quality assessed using the Scottish Intercollegiate Guidelines Network methodology checklist for cohort studies. The search revealed 898 articles narrowed to full-text review of 23 studies. In total, 11 research articles were included in the review, describing nine different studies. The findings were narratively summarized in line with the aims of the review.

Findings

The included studies showed marked heterogeneity, which impacts on the findings. However, the majority report an increase in energy intake (kcal or kJ) during pregnancy. Of the studies that reported changes through food group comparisons, a majority reported a significant increase in fruit and vegetable consumption, a decrease in egg consumption, a decrease in fried and fast food consumption and a decrease in coffee and tea consumption from before to during pregnancy. The characteristics of the women participating in these studies, suggest that age, education and pregnancy intention are associated with healthier dietary changes; however these factors were only assessed in a small number of studies.

Key conclusions

The 11 included articles show varied results in dietary intake during pregnancy as compared to before. More research is needed regarding who makes these healthy changes, this includes consistency regarding measurement tools, outcomes and time points.

Implications for practice

Midwives as well as intervention developers need to be aware of the dietary changes women may spontaneously engage in when becoming pregnant, so that care and interventions can build on these.

Keywords: Pre-conception, pregnancy, dietary intake, caffeine, systematic review.

ACCEPTED MANUSCRIPT

Introduction

Pregnancy is a time when many women gain weight they subsequently retain post pregnancy. Almost 30% of pregnant women gain more weight than is recommended by the American Institute of Medicine guidelines (IOM, 2009) and previous evidence from a range of countries suggests a mean weight gain between 0.4kg to 3.8 Kg as a result of pregnancy up to 2.5 years postpartum (Linne et al., 2002). Increased weight post-natal was also found to be a strong indicator of being overweight 15 years later (Linne et al., 2004). The more weight gained during pregnancy, the more likely that it may be retained postpartum (Johnson et al., 2013) and women who enter a subsequent pregnancy overweight or obese also have a higher risk of adverse outcomes for themselves and/or their infants (Kuhlmann et al., 2008 and Marchi et al., 2015).

Numerous interventions have targeted weight gain in pregnancy, including both physical activity and dietary components. A recent review suggests that interventions with dietary aspects may be most effective in helping women gain a healthy weight in pregnancy (Thangaratinam et al., 2012). Adequate nutritional intake during pregnancy is vitally important to ensure appropriate fetal growth both physically and mentally (Anderson et al., 2001) and poor maternal nutritional status is well reported to not only affect pregnancy outcomes (Osrin and de L Costello, 2000 and Keen et al., 2003), but may also be related to the risk of developing several non-communicable diseases in the adult child (Barker et al, 2013). As such dietary intake both before and during pregnancy is a major public health issue (Barker et al., 2013).

Pregnancy is a period where women are particularly concerned with their dietary intake (Pinto et al., 2008) and are considered highly motivated for dietary improvements (Szwajcer et al., 2008 and Phelan, 2010). For example, when pregnant women have been asked for the behaviours they do to keep healthy in pregnancy, healthy eating is the most commonly mentioned health behaviour (Lewallen, 2004). To ensure appropriate and effective interventions are successfully developed to improve dietary intake during pregnancy, the dietary changes women make when they become pregnant are important to understand (Skreden et al., 2014). The primary aim of this systematic review was therefore to review the existing literature on dietary intake change before and during pregnancy. In addition to knowing *what* dietary changes women make when becoming pregnant, it is also important to understand *who* makes these changes. Thus, our secondary review aim was to identify the key characteristics of the women who report changing their dietary intake from before to during pregnancy.

Methods

A systematic literature review was conducted to identify the changes in women's dietary intake before and during pregnancy and to identify which women may make these changes. Three databases (Web of Science, CINAHL and PubMed) were systematically searched in May, 2016. Search terms included preconception, pre-pregnancy, pregnancy, gestation, dietary intake, food intake, beverages, caffeine, fruit and vegetables. Scopus was used for forward searching (May 2016). Studies were included if they measured women's dietary intake before and during pregnancy, either prospectively or retrospectively. For the purpose of this review, dietary intake included food groups as well as energy and macronutrients. Notably, drinking alcohol was not included in this review despite being part of a woman's energy intake. There are two reasons for omitting alcohol from this review, firstly not all women drink alcohol when not pregnant (Petherick et al., 2010). Secondly, drinking alcohol is consistently reported to decrease before and during pregnancy (Crozier et al., 2009a; Aden et al., 2007; Pinto et al., 2008).

In addition, to be included studies had to use a within-participants design to limit the bias and individual variance associated with dietary intake. Lastly, to be included, articles had to be in English and in peer-reviewed journals. Screening of titles and abstracts and decision on final inclusion of articles was done by both authors.

Analysis

All papers were quality assessed using the Scottish Intercollegiate Guidelines Network methodology checklist for cohort studies (Scottish Intercollegiate Guidelines Network, 2016). This checklist was chosen as it differentiates between prospective and retrospective cohort studies, of which both were included in this review. Both authors scored the studies independently and scoring discrepancies were resolved via discussion. Inter-rater reliability was calculated using percentage agreement.

For all studies, study population, study design, diet measurement, type and timing of measurement and study findings were extracted. Due to the heterogeneity of the studies identified it was inappropriate to conduct a meta-analysis and a narrative method of synthesis analysis was conducted. This method has been used previously when the experimental studies included are not sufficiently similar for a meta-analysis to be appropriate (Mays et al., 2005) Ethical approval was not required for this systematic review.

Results

The literature search yielded 898 articles including one article found by a hand search, of which 468 were screened by title and abstract and 23 were full text screened (see Figure 1). Details of study exclusion are detailed in Table 2 in supplementary material. Forward searching identified two additional articles (Aden et al., 2007 and Crozier et al., 2009a). In total, 11 research articles were included in the review, describing nine different studies.

Study characteristics

The included studies heralded from all over the world, published between 1998 and 2014 (see Table 1). The majority of studies used a prospective design (n= 6) with three studies (reported in five articles) using a retrospective design. Study sample size varied from 10 (Kopp-Hoolihan et al., 1999) to 7174 (Hellerstedt et al., 1998). The included studies varied greatly regarding the information authors reported regarding participant characteristics in terms of age, ethnicity, parity and weight status (see Table 1). Variations in measurement time points were also noticed with the prospective studies measuring pre-conception dietary intake within a few months of a confirmed pregnancy. The retrospective studies measured dietary intake at different time points during pregnancy or postpartum to gather information of dietary intake before and during pregnancy. Dietary intake was either measured through interview (face-to-face or by phone) or self-administered questionnaire. In total, seven articles provided data on changes in food groups and three articles reported findings in terms of energy and macronutrients, with one reporting both methods. Four articles provided data on characteristics of the women who report changing their dietary intake before and during pregnancy.

Quality assessment

Inter-rater reliability, assessed through percentage agreement was 77.8%. Whilst the prospective studies were deemed marginally stronger compared to the retrospective studies, all articles were found to be of acceptable quality. See Table 1 in supplementary material for full breakdown of quality assessment.

The results of the review are presented under two headings, dietary intake changes and characteristics of women making dietary changes. Changes in dietary intake will be clustered using the sub-headings of food groups or energy and macronutrient intake to complement the individual study reporting and to allow comparisons between studies to be made more easily.

Dietary intake change from preconception to pregnancy

Food Groups - Fruits and vegetables

Six articles reported data on fruit and vegetable intake with inconsistent findings (Cuco et al., 2006a; Pinto et al., 2008; Crozier et al., 2009a; Crozier et al., 2009b; Paulik et al., 2009; Smedley et al., 2014). Paulik et al. (2009) reported an increase in the percentage of women consuming both fruits and vegetables (more than 4 times per week) in pregnancy (85.7% vs 94.8% fruit and 67.6% vs 75.4% vegetables). This is further supported by Smedley et al. (2014), who reports a significant increase in the number of women 'always' consuming fruit and vegetables during pregnancy (65% vs 78% fruit and 61% vs 77% vegetables). Crozier et al. (2009a) reports an increase in citrus fruit and fruit juice intake during pregnancy compared to before pregnancy (52% vs. 64%). In contrast, Pinto et al. (2008) reported no significant change in median daily vegetable consumption (grams) between preconception and pregnancy, but did report a significant increase in fruit consumption during pregnancy (+21.5 grams). This was also supported by Cuco et al. (2006a) who reports no significant differences in mean consumption of fruit or vegetable intakes. In addition, portions of fruit and vegetables per day did not significantly differ between pre-conception and during pregnancy (5.2 vs. 5.35 portions) as reported by Crozier et al., (2009b).

Dairy

Three studies reported data on dairy intake and the results varied greatly between studies (Pinto et al., 2008; Crozier et al., 2009a; Smedley et al., 2014). Pinto et al. (2008) reported a significant increase in milk and dairy products between pre-conception and during pregnancy (387.5g vs 691.8g), and a significant decrease in egg consumption between pre-conception and during pregnancy (22.2g vs 11.1g). In addition, Crozier et al. (2009a) reported an increased intake in a number of dairy products including cream and milk as well as reporting an increase in the consumption of cheese and cottage cheese during both early (3.0 portions) and late (4.5 portions) pregnancy when compare to pre-conception (1.8 portions). However Smedley et al. (2014) reported no significant difference in dairy intake in all categories between pre-conception and during pregnancy

Meat and meat products

Two studies reported data on meat and meat products and the results varied greatly between studies (Pinto et al., 2008; Crozier et al., 2009a). Crozier et al. (2009a) reported an increase in processed meat consumption during early and late pregnancy, but reported no change in red meat, chicken, turkey or fish consumption during pregnancy. Crozier et al. (2009a) also reported that the proportion of women consuming meat such as liver and kidneys was 48% during pre-conception,

and decreased to 22% in early pregnancy and 16% in late pregnancy. This contrasts with evidence reported by Pinto et al. (2008) who reported a significant decrease in red meat consumption during pregnancy (-4.7g) but who also found no significant difference in fish consumption.

Starchy Carbohydrates (CHO)

Two studies reported data on starchy carbohydrates and the results varied greatly between studies (Pinto et al., 2008, Crozier et al., 2009a). Pinto et al. (2008) reported a significant increase in bread consumption but a decrease in rice, pasta and potato consumption during pregnancy. Crozier et al. (2009a) reported that rice and pasta consumption was lower during early and late pregnancy with an increase in weekly consumption of breakfast cereals during late pregnancy (7 portions) compared to pre-conception (4.5 portions) and early pregnancy (4.5 portions) also reported. However Crozier et al. (2009a) also reported no changes in intake of wholemeal bread, quiche, pizza and pancakes.

Sweet foods

Three studies reported data on sweet foods and the results varied greatly between studies (Pinto et al., 2008; Crozier et al., 2009a; Smedley et al., 2014). Smedley et al. (2014) and Pinto et al. (2008) reported no change in sweet bakery food or sweets consumption between pre-conception and during pregnancy, whereas Crozier et al. (2009a) reported an increase in portion consumption of sweet spreads, confectionary, cakes and biscuits during both early and late pregnancy, whereas puddings only increased during late pregnancy.

Fast and Fried Food

Two articles reported data on fried and fast food (Pinto et al, 2008; Smedley et al, 2014). Fried food intake was not significantly different before and during pregnancy (Smedley et al., 2014). However fast food intake did decrease during pregnancy, with a greater number of women reporting that they 'never' consumed this food (56% vs 67%) (Smedley et al., 2014). Similarly, Pinto et al. (2008) reported a decrease in the consumption of fast food during pregnancy compared to pre-conception intake (25.1g vs 17.1g)

Beverages

Five articles reported data on beverage intake and the results varied greatly between studies (Hellerstedt et al., 1998; Cuco et al. 2006a; Pinto et al., 2008; Crozier et al., 2009b; Skreden et al., 2014). Coffee and tea was the most commonly reported beverage, and consumption was found to decrease from before to during pregnancy in four studies (Hellerstedt et al., 1998; Pinto et al., 2008;

Crozier et al., 2009b; Skreden et al., 2014). Paulik et al. (2009, n=349) reports a decrease in drinking one cup of coffee per day from 56.2% to 33.2%. Milk was assessed in three articles; Skreden et al. (2014) reported an increase in milk intake and Pinto et al. (2008) reported an increase in daily intake of milk and dairy products. Whereas Paulik et al. (2009) reported a decrease from 66.8% vs 60.2%.

Regarding sugar sweetened beverages and fruit juices, both Pinto et al. (2008) and Cuco et al. (2006a) reported no significant differences from before to during pregnancy, whilst Crozier et al. (2009a) and Skreden et al. (2014) found an increase in fruit juice consumption. Moreover, a decrease in sugar-sweetened beverages and artificially sweetened beverages was found in both studies (Crozier et al., 2009a and Skreden et al., 2014). Cuco et al. (2006a) also reported that participants who had high scores of sweetened beverages and sugar during both pre-conception and during pregnancy tended to consume less fresh fruit, vegetables, roots and tubers. Lastly, the percentage of women who reported at least daily consumption of water increased from before to during pregnancy (Skreden et al., 2014).

Energy and Macronutrients

Total energy intake (kcal, kJ or MJ) was measured in five studies (Koop-Hoolihan, 1999; Pinto et al., 2008; Cuco et al., 2006a and 2006b and Aden et al., 2007), with four studies (Koop-Hoolihan, 1999; Cuco et al., 2006a and 2006b and Aden et al., 2007) recording an increase in energy intake during pregnancy and one reporting no significant change (Pinto et al., 2008).

Kopp-Hoolihan et al. (1999) reported energy intake using three day food diaries from 10 women during pre-conception (T1) and three trimesters during pregnancy (T1, T2, and T3). The results show a 9% increase (775kJ/day) in total energy intake between T1 and T3. Similarly, Aden et al. (2007) reported an increase in energy intake between pre-conception (1852 ± 751 kcal/day) and during pregnancy (2104 ± 583 kcal/day) using a Food Frequency Questionnaire (FFQ) and a 24hr dietary recall, although there was no indication if this was a statistically significant increase. Cuco et al. (2006a and 2006b) reported changes in energy intake between pre-conception and four different weeks during pregnancy. The authors in both articles report an increase in energy intake between preconception and the 10th and 26th week of pregnancy but a decrease during the 6th and 38th week. However, Pinto et al. (2008) reported no significantly difference between pre-conception (2393 kcal/day) and during pregnancy (2423 kcal/day).

Macronutrient intake was also reported in 3 studies (Cuco et al., 2006b; Aden et al., 2007 and Pinto et al., 2008), with no consistent changes in intake reported in studies. Cuco et al. (2006b) reported macronutrient intake using a 7 day consecutive food diary. Protein intake did not differ between pre-conception and during pregnancy; however the proportion of animal to vegetable protein increased in favour of vegetable protein during pregnancy compared to pre-conception. CHO and fat intake increased during the 10th, 26th and 38th week (182.2g preconception vs 199.4g; 206.7g; 191.8g respectively CHO and 91.6g preconception vs 98.0g, 97.3g, 92.9g respectively Fat). Cuco et al. (2006b) also reported changes in maternal consumption of protein, fat, CHO and suggests that an increase of only 1 gram of these during preconception, 6th, 10th, 26th and 38th week of pregnancy can cause significant changes in child birth weight (7.8 – 11.4 grams)

Aden et al. (2007) reported an increase in CHO and protein intake with a decrease in fat intake recorded during pregnancy. However, Pinto et al. (2008) reported no significant differences between CHO and total fat intake as a percentage of total energy intakes (%TEI) between pre-conception and during pregnancy. However the results do indicate a significant increase in %TEI saturated fat (SFA) and protein during pregnancy compared to pre-conception.

Characteristics of the women who report changing their dietary intake from before to during pregnancy

Four studies reported characteristics of the women who made dietary changes from before to during pregnancy. Crozier et al. (2009b) explored what variables may predict daily fruit and vegetable intake. They found that both at pre-conception and during pregnancy, younger women ate less than five portions of fruit and vegetables a day compared to older women. Cuco et al. (2006a) also reports a positive association between the consumption of vegetables and meat with age. Whilst Skreden et al. (2014) found that women over 25 years reported larger decreases in artificially sweetened beverages and increased their fruit juice consumption more compared to women less than 25 years old. The women over 25 years also reported a larger intake in milk compared to younger women from pre-conception to during pregnancy. The same study found no relationship between pre-pregnancy BMI (>25 vs. <25) and changes in drinking habits or beverage consumption. Skreden and colleagues (2014) also found that higher education was associated with more reduction in coffee consumption. Lastly, Hellerstedt et al. (1998) examined daily caffeine use and pregnancy intention. They found that women with intended pregnancies, compared to those who reported the pregnancy was unintended, were more likely to report decreased consumption of caffeine from before to during pregnancy.

Discussion

The aims of this review were to evaluate the evidence relating to what changes in dietary intake women make when becoming pregnant, and secondly identify any characteristics of the women making these changes. The included studies are heterogeneous, specifically in relation to outcome measures and time frames in which data collection occurred; as such the findings should be interpreted with caution. Overall, the review findings suggest that some changes regarding dietary intake are made during pregnancy and these are in line with studies that have compared dietary intake between pregnant women and non-pregnant women (Anderson et al, 1993; Verbeke et al, 2007 and Inskip et al, 2009). The majority of studies report an increase in energy intake (kcal or kJ) during pregnancy, but failed to consistently report changes in different macronutrient intake (Cuco et al., 2006a and 2006b; Aden et al., 2007 and Pinto et al., 2008). Of the studies that reported changes through food group comparisons, a majority reported a significant increase in fruit and vegetable consumption, a decrease in egg consumption, a decrease in fried and fast food consumption and a decrease in coffee and tea consumption from pre-conception to during pregnancy (Helderstedt et al., 1998; Cuco et al., 2006a; Pinto et al., 2008; Crozier et al., 2009a and 2009b; Paulik et al., 2009; Skreden et al., 2014; Smedley et al., 2014). There was no consistency in starch carbohydrate consumption, meat, fish or sweets/sweet food consumption. Regarding the characteristics of the women making these dietary changes, only three studies provided information and as such no conclusions can be drawn.

Dietary intake change before and during pregnancy

Changes in energy intake were found to vary considerably between studies, with several papers reporting a significant increase and others reporting no significant change. Despite the general trend towards an increase in overall energy intake there were no consistent differences reported in specific macronutrient intake from before and during pregnancy. However one author (Aden et al., 2007) did report a large range in energy intake between both stages, with pre-conception intake ranging between 1116 kcal/day to 6087 kcal/day and during pregnancy ranging between 945 kcal/day and 3627 kcal/day. This indicates that although average intake may not change, there are likely to be large inter-individual variations in the overall energy and macronutrient intake between pregnant women which could have significant health and weight implications.

In terms of food group consumption, the most consistent findings are an increase in fruit and vegetable intake as well as an increase in dairy and a decrease in caffeine intake. An increase in fruit intake has also been reported in studies comparing pregnant to non-pregnant women (Anderson et

al, 1993; Verbeke et al, 2007), although one study found little difference between these groups (Inskip et al., 2009). Although fruit and vegetable intake was widely reported to increase during pregnancy, it cannot be assumed that all women adequately consumed the national recommendations for fruit and vegetable consumption per day. Smedley et al. (2014) reported that although fruit and vegetable consumption increased during pregnancy, only two thirds of participants reported consuming the recommend quantities of fruit and vegetable as suggested by the Australian public health guidelines (National Health and Medical Research Council, 2003). As fruit and vegetable intake is recommended as part of a healthy balanced diet, and their increased consumption is linked with a number of positive health outcomes (Slavin and Lloyd, 2012), the results indicate that more information should be provided to women before and during pregnancy on the importance of not only increased fruit and vegetable consumption but to ensure they reach the correct public health recommendations for their country.

Two studies found an increase in milk and dairy consumption (Pinto et al. 2008 and Crozier et al., 2009a). This is in line with other research findings where pregnant women report higher dairy intake compared to non-pregnant women (Anderson et al, 1993; Verbeke et al, 2007). This increase is positive as the recommended intake of calcium increases during pregnancy and studies reporting micronutrient intake only indicate that calcium intake increases during pregnancy (Aden et al., 2007) which could further explain the reported increase in dairy consumption (Crozier et al., 2009a and Pinto et al., 2008). The increase in dairy consumption could also account for the increase in energy intake recorded (Koop-Hoolihan, 1999; Cuco et al., 2006a and 2006b and Aden et al., 2007), particularly as the types of products consumed may correspond to more energy-dense foods such as full-fat milk and cheese (Crozier et al., 2009a).

In terms of beverages, there was encouraging findings that women decrease their coffee intake when pregnant and increase their milk intake. A decrease in daily caffeine intake has also been found in women attempting pregnancy (Lum et al., 2011), this suggests it is a component of healthy eating some women are aware of. In terms of fruit juices and sugar-sweetened drinks, two studies reported inconsistent findings, and more research is needed. Fruit juices and sugar-sweetened drinks are both important to target for weight-management as they are often high in calories.

In addition, the proportion of women consuming liver and kidneys was 48% pre-conception, 22% in early pregnancy and 16% in late pregnancy (Crozier et al., 2009a); this change in consumption is consistent with previous public health messages in pregnancy relating to the harmful effects of

excess vitamin A consumption through liver consumption (NHS Choices, 2015), despite little scientific evidence to support this (Strobel et al., 2007). Similarly the decrease of consumption in fast food reported (Smedley et al., 2014) could be due to public health education programmes in Australia relating to foods not to eat to avoid Listeria (Anderson, 2001). Indeed, previous research has suggested that health education around effective weight management can affect weight gain during pregnancy (Wilkinson et al., 2009), with further evidence to suggest that pre-conception interventions can improve both the intention and self-efficacy of healthy eating behaviours during pregnancy (Hillemeier et al., 2008). There is also emerging evidence to suggest that women start eating healthily in preparation for pregnancy (Ramage et al, 2015).

The variation in dietary intake changes reported before and during pregnancy in the reviewed studies, may be due to the disparity of nutritional and lifestyle advice given by different countries (Shawe et al., 2015). A recent publication by Shawe et al. (2015) reviewed the pre-conception care policy, guidelines and recommendations of six European countries (Belgium, Denmark, Italy, Netherlands, Sweden and UK) and reported that there were large variations between countries particularly in relation to fish, caffeine and alcohol consumption. This could account for some of the inconsistent results reported by the current studies reviewed.

Characteristics of the women who report changing their dietary intake from before to during pregnancy

Only four studies reported characteristics of the women making dietary changes. Findings suggest that education and age may be linked to dietary intake (Crozier et al. 2009b; Cuco et al. 2006a; Skreden et al., 2014) where older and more educated women tend to make healthier dietary changes. Findings from one study suggest that pregnancy intention may be associated with coffee intake (Hellerstedt et al., 1998). Since our search, a recent study fitting the scope of our review has been published where older pregnant women were more likely to decrease their intake of processed foods compared to younger pregnant women (Alves-Santos et al, 2016). Thus, whilst it is disappointing that so few studies examined the demographic and pregnancy factors that may be associated with dietary changes, our findings suggest that age, education and pregnancy intention may be factors worthy further examination. For example, nutrition awareness has been found to be higher in women trying to conceive compared to those women not trying to conceive (Szwajcer et al, 2012). This information is likely to be important for targeting the right population of women with interventions and support.

Strengths and limitations

There are a number of strengths and limitations relating to the evidence presented in this review. Quality assessment of the 11 studies included using the SIGN checklist, reported the studies to be acceptable or highly acceptable in quality (Scottish Intercollegiate Guidelines Network, 2016). This indicates that despite the relatively low number of articles meeting the inclusion criteria (n=11) they were overall of good quality. Another strength was the range of countries in which the data was collected from, showing consistency in dietary change across different cultures although only English language articles were included.

One limitation of the literature included in the review is the different methods used to measure dietary intake. Ranging from food frequency questionnaires (FFQ), food diaries (FD; 3 and 7 day; weighed and unweighed) as well 24 hour dietary recall methods. Pinto et al. (2008) justified the use of an FFQ in their study as it allowed for retrospective estimation of dietary intake to be collected. However they also recorded intake with a 3 day food diary during pregnancy (Pinto et al., 2008) and reported that differences in intake recorded between the methods may be due to previous evidence indicating that the FFQ tends to overestimate intake whereas FD tends to underestimate (Cade et al., 2002). In addition, the longer the period of dietary recording, the greater likelihood of participant fatigue and therefore potential under or overestimation of dietary intake (Buzzard, 1998).

Studies included in this review were both prospective and retrospective in nature. Retrospective studies are limited in quality as they are subject to participant recall bias and potentially the prior knowledge of pregnancy outcomes could have affected the outcome of dietary intake recall (Pinto et al., 2008). In addition, recall bias may have been greater in women who experienced nausea and vomiting in early pregnancy and this may have affected dietary intake patterns when comparing pre-conception to during pregnancy (Pinto et al., 2008). Furthermore, the diversity of time points used by researchers is problematic, as women may change their eating throughout pregnancy. That said, those studies that measured diet at different time points in pregnancy report inconsistent findings regarding whether diet changes or not (Pinto et al., 2008; Cuco et al., 2006a). Clearly more research is needed. Not all included papers in this review reported changes in dietary intake as a primary objective and thus not conducting significance testing. These papers were still included due the authors to wanting to include all identified evidence in the review.

In addition, this review only included studies if they used a within-group study design. It must be acknowledged that studies using this design are subject to a number of limitations including practice

effects and fatigue, with participants potentially becoming more attuned to detailing their dietary intake practices, increasing the likelihood of miss-reporting. As such, this needs to be considered when interpreting the results. It must also be acknowledged that the review question could have been answered using other research designs such as comparisons between groups of pregnant and non-pregnant women. We have compared our review findings with evidence from such studies in the Discussion section, and shown that our findings are in line with these studies.

Implications and future directions

This review provides implications for both healthcare professionals, such as midwives, and intervention developers. Women often report wanting information early in their pregnancy about healthy eating (Olander et al., 2012). Healthcare professionals are consistently identified as the key source of information regarding healthy diet in pregnancy (Olander et al., 2012 and Smedley et al., 2014) and thus it is important for midwives and others to be aware of the dietary changes women may make when becoming pregnant, so that positive changes can be supported and built upon. It is also important to be mindful that a planned pregnancy may not necessarily mean women are healthier in preconception, and thus are likely to need the same advice as those women who have an unplanned pregnancy.

For intervention developers, these review findings are important to consider when targeting dietary intake in pregnancy. This review has identified food groups and characteristics of women that may confound intervention results. The review identifies that future studies should develop an agreed set of measures (timeframes, dietary recording techniques) for use across studies on this topic to reduce the problem of heterogeneity in this area. A successful intervention must be able to identify what behaviours women may change spontaneously when becoming pregnant and what behaviours they need support with.

Conclusion

Dietary intake before and during pregnancy has significant implications for the mother and unborn child with a number of health outcomes related to poor dietary intake. The current literature available on women's change in dietary intake, using within-subject design, from before to during pregnancy is limited to a handful of studies using a variety of dietary intake recording methods on a wide range of dietary variables to collect data both prospectively and retrospectively and whose overall quality is acceptable or highly acceptable. The evidence suggests that a number of changes in dietary intake may take place during pregnancy (such as an increase in fruit and vegetable intake),

464 but that a number of other key components relating to high energy dense foods are inconsistent
465 which could have far reaching implications in terms of energy balance and excess weight gain during
466 pregnancy. Further research needs to be conducted investigating the changes in dietary intake
467 before and during pregnancy prospectively, using this alongside records of weight gain and
468 pregnancy outcomes in both mother and child to determine the longer term health implications of
469 poor dietary intake.

ACCEPTED MANUSCRIPT

References

- Aden, E., Johansson, I. and Haglin, L. 2007. Energy and nutrients in self-reported diet before and at week 18-22 of pregnancy. *Scandinavian Journal of Food and Nutrition* 51(2), 67-73
- Anderson, A.S., Campbell, D., and Shepherd, R. 1993. Nutrition knowledge, attitude to healthier eating and dietary intake in pregnant compared to non-pregnant women. *Journal of Human Nutrition and Dietetics*, 6(4), 335-353
- Anderson, A.S. 2001. Pregnancy as a time for dietary change? *Proceedings of the Nutrition Society* 60, 497-504
- Alves-Santos, N.H., Eshriqui, I., Franco-Sena, A.B., Cocate, P G., Freitas-Vilela, A.A., Benaim, C., Vaz Jdos, S., Castro. M.B. and Kac, G. 2016. Dietary intake variations from pre-conception to gestational period according to the degree of industrial processing: A Brazilian cohort. *Appetite*, 105, 164-171.
- Barker, D., Barker, M., Fleming, T. 2013. Developmental biology: support mothers to secure public health. *Nature* 504, 209-211
- Buzzard, M. 1998. 24-hour dietary recall and food record methods. In: Willet W, editor. *Nutritional Epidemiology*, 2nd edn. New York: Oxford University Press, 50-73
- Cade, J., Thompson, R., Burley, V. and Warm, D. 2002. Development, validation and utilisation of food-frequency questionnaires – a review. *Public Health Nutrition* 5, 567-587
- Crozier, S., Robinson, S., Godfrey, K., Cooper, C. and Inskip, H. 2009a. Women's dietary patterns change little from before to during pregnancy. *The Journal of Nutrition* 139, 1566-1573
- Crozier, S., Robinson, S., Borland, S., Godfrey, K., Cooper, C., Inskip, H. and the SWS Study Group. 2009b. Do women change their health behaviours in pregnancy? Findings from the Southampton Women's Survey. *Paediatric and Perinatal Epidemiology* 23, 445-453.
- Cuco, G., Fernandez-Ballart, J., Sala, J., Viladrich, C., Iranzo, R., Vila, J. and Arija, V. 2006a. Dietary patterns and associated lifestyles in preconception, pregnancy and postpartum. *European Journal of Clinical Nutrition* 60, 364-371
- Cuco, G., Arija, V., Iranzo, R., Vila, J., Prieto, M. and Fernandez-Ballart, J. 2006b. Association of maternal protein intake before conception and throughout pregnancy with birth weight. *Acta Obstetrica et Gynecologica*. 85, 413-421

- 498 Hellerstedt, W., Pirie, P., Lando, H., Curry, S., McBride, C., Grothaus, L., and Nelson, J. 1998.
499 Differences in preconceptional and prenatal behaviours in women with intended and unintended
500 pregnancies. *American Journal of Public Health* 88(4), 663-666.
- 501 Hillemeier, M., Symons Downs, D., Feinbery, M., Weisman, C. S., Chuang, C., Parrott, R., Velott, D.,
502 Francis, L., Baker, S., Dyer, AM. Chinchilli, V. 2008. Improving Women's Preconceptional Health:
503 Findings from a Randomized Trial of the Strong Healthy Women Intervention in the Central
504 Pennsylvania Women's Health Study. *Women's Health Issues* 18, (6), S87-S96
- 505 Inskip, H.M., Crozier, S.R., Godfrey, K.M., Borland, S.E., Cooper, C., and Robinson, S.M. 2009.
506 Women's compliance with nutrition and lifestyle recommendations before pregnancy: general
507 population cohort study. *British Medical Journal* 338, b481
- 508 Institute of Medicine. 2009. Weight gain during pregnancy: re-examining the guidelines.
509 Washington, DC: National Academies Press
- 510 Johnson, M., Campbell, F., Messina, J., Preston, L., Buckley Woods, H. and Goyder, E. 2013. Weight
511 management during pregnancy: A systematic review of qualitative evidence. *Midwifery* 29(12),
512 1287-96.
- 513 Keen, C., Clegg, M., Hanna, L., Lanoue, L., Rogers, J. and Daston, G. 2003. The plausibility of
514 micronutrient deficiencies being a significant contributing factor to the occurrence of pregnancy
515 complications. *Journal of Nutrition* 133(Suppl 2), S1597-S1605
- 516 Kopp-Hoolihan, L., van Loan, M., Wong, W. and King, J. 1999. Longitudinal assessment of energy
517 balance in well-nourished, pregnant women. *American Journal of Clinical Nutrition* 69, 697-704
- 518 Kuhlmann, A.K., Dietz, P.M., Galavotti, C. and England, L.J. 2008. Weight-management interventions
519 for pregnant or postpartum women. *American Journal of Preventive Medicine* 34(6), 523-8
- 520 Lewallen, L.P. 2004. Healthy Behaviors and Sources of Health Information Among Low-Income
521 Pregnant Women. *Public Health Nursing* 21(3), 200-6.
- 522 Linne, Y., Barkeling, B., Rossner, S. 2002. Long-term weight development after pregnancy. *Obesity*
523 *Reviews*. 3, 75-83.
- 524 Linne, Y., Dye, L., Barkeling, B. and Rossner, S. 2004. Long-term weight development in women: a 15-
525 year follow-up of the effects of pregnancy. *Obesity Research* 12(7), 1166-78

- 526 Lum, K. J., Sundaram, R. and Buck Louis, G.M. 2011. Women's lifestyle behaviours while trying to
527 become pregnant: Evidence supporting preconception guidance. *American Journal of Obstetrics and*
528 *Gynecology* 205(3), e201-203.
- 529 Marchi, J., Berg, M., Dencker, A., Olander, E. and Begley, C. 2015. Risks associated with obesity in
530 pregnancy, for the mother and baby: a systematic review of reviews. *Obesity Reviews* 16(8), 621-
531 638.
- 532 Mays, N., Pope, C., and Popay, J., 2005. Systematically reviewing qualitative and quantitative
533 evidence to inform management and policy-making in the health field. *Journal of Health Service*
534 *Research and Policy*, 10 (Suppl 1), 6–20.
- 535 National Health and Medical Research Council. Food for health – Dietary guidelines for Australians.
536 Department of Health and Aging (ed.). Canberra: Commonwealth of Australia, 2003
- 537 NHS Choices (2015) Why should I avoid some foods during Pregnancy?
538 <http://www.nhs.uk/chq/Pages/917.aspx?CategoryID=54>. Accessed [31.08.16]
- 539 Olander, E.K., Atkinson, L., Edmunds, J.K. and French, D.P. 2012. Promoting healthy eating in
540 pregnancy: What kind of support services do women say they want? *Primary Health Care Research*
541 *and Development* 13, 237-243.
- 542 Osrin, D., de L Costello, A. 2000. Maternal nutrition and fetal growth: practical issues in international
543 health. *Seminar Neonatal* 5, 209-219
- 544 Paulik, E., Csaszar, J., Kozinszky, Z. and Nagymajtenyi, L. 2009. Preconceptual and prenatal predictors
545 of folic acid intake in Hungarian pregnant women. *European Journal of Obstetrics and Gynecology*
546 *and Reproductive Biology* 145, 49-52
- 547 Petherick, E., Parslow, R., McKinner, P., Tufnell, D., Leon, D., Raynore, P., Lawlor, D. and Wright, J.
548 2010. P13 Association of prenatal and postnatal smoking and alcohol consumption on birth weight in
549 the white British population in Bradford: preliminary findings from the born in Bradford study.
550 *Journal of Epidemiology & Community Health* 64, A38-39 31p.
- 551 Phelan, S. 2010. Pregnancy: a 'teachable moment' for weight control and obesity prevention.
552 *American Journal of Obstetrics and Gynaecology* 202 (135), e131-e138
- 553 Pinto, E., Barros, H., and Santos Silva, I. 2008. Dietary intake and nutritional adequacy prior to
554 conception and during pregnancy: a follow-up study in the north of Portugal. *Public Health Nutrition*
555 12(7), 922-931

- 556 Ramage, S.M., McCargar, L.J., Berglund, C., Harber, V., and Bell, R.C. 2015. Assessment of pre-
557 pregnancy dietary intake with a food frequency questionnaire in Alberta women. *Nutrients* 7(8),
558 6155-6166.
- 559 Scottish Intercollegiate Guidelines Network. <http://www.sign.ac.uk/methodology/checklists.html>
560 Accessed 27.5.16
- 561 Shawe, J., Delbaere, I. Ekstrand, M. Hegaard, H. K. Larsson, M. Mastroiacovo, P., Stern, J., Steegers,
562 E., Stephenson, J. and Tyden, T. (2015). Preconception care policy, guidelines, recommendations and
563 services across six European countries: Belgium (Flanders), Denmark, Italy, the Netherlands, Sweden
564 and the United Kingdom. *European Journal of Contraception & Reproductive Health Care* 20(2), 77-
565 87
- 566 Skreden, M., Bere, E., Sagedal, L.R., Vistad, I., Øverby, N.C. 2014. Changes in beverage consumption
567 from pre-pregnancy to early pregnancy in the Norwegian Fit for Delivery study. *Public Health*
568 *Nutrition* 18 (7), 1187-1196 .
- 569 Slavin, J. L. and Lloyd, B. 2012. Health benefits of fruits and vegetables. *Advances in Nutrition* 3, 506-
570 516.
- 571 Smedley, J., Jancey, J., Dhaliwal, S., Zhao, Y., Monteiro, S. and Howat, P. 2014. Women's reported
572 health behaviours before and during pregnancy; A retrospective study. *Health Education Journal* 73
573 (1), 28-40
- 574 Strobel, M., Tinz, J and Biesalski, H.K. 2007. The importance of beta-carotene as a source of vitamin
575 A with special regard to pregnant and breastfeeding women. *European Journal of Nutrition*. 46,
576 Suppl.1, I1-20
- 577 Szwajcer, E., Hiddink, G. and Maas, L. 2008. Nutrition-related information-seeking behaviours of
578 women trying to conceive and pregnant women: evidence for the life course perspective. *Family*
579 *Practice* 25(Suppl. 1), i99-i104
- 580 Szwajcer, E., Hiddink, G.J., Maas, L., Koelen, M., and van Woerkum, C. 2012. Nutrition awareness
581 before and throughout different trimesters in pregnancy: a quantitative study among Dutch
582 women. *Family Practice* 29 (Suppl 1), i82-i88.
- 583 Thangaratinam, S., Rogozinska, E., Jolly, K., Glinkowski, S., Roseboom, T., Tomlinson, J. W. and Khan,
584 K. S. 2012. Effects of interventions in pregnancy on maternal weight and obstetric outcomes: meta-
585 analysis of randomised evidence. *British Medical Journal* 344, e2088.

586 Verbeke, W. and De Bourdeaudhuij, I. 2007. Dietary behaviour of pregnant versus non-pregnant
587 women. *Appetite* 48(1), 78-86.

588 Wilkinson, S., Miller, Y. and Watson, B. 2009. Prevalence of health behaviours in pregnancy at
589 service entry in a Queensland health service district. *Australian and New Zealand Journal of Public*
590 *Health* 33(3), 229-23.

591


ACCEPTED MANUSCRIPT

592 **Acknowledgements:**

593 We are very grateful for the comments made by two anonymous reviewers on our submitted
594 manuscript. These comments helped strengthen the manuscript.

ACCEPTED MANUSCRIPT

Figure 1. Flowchart describing the number of articles retrieved, and included and excluded at each stage of the review process



598 **Table 1.** Summary of studies included in review.

Study authors, year (country)	Sample characteristics	Study design (prospective / retrospective)	When was diet measured?	What was measured?	How was diet measured?	Change in diet	Women's characteristics
Aden et al, 2007 (Sweden)	Sample size: 50 Age: 30 years (SD 4.6; range 18-40 years) Gestation weeks: 18.1 (SD 1.1; range 15-21) weeks Weight category: Pre-pregnant BMI mean 23.2 (SD3.1, 17.1-32.4) SES or similar: Before pregnancy FT working 52% Student 20% PT working 16% Unemployed/sick leave 4% Other 8% Education: Not reported Ethnicity: Swedish 92% Asian 6%, Persian 2% Smoking: Pre-pregnancy 16%, during pregnancy 6% NCD's: Not reported Parity: 62% first time mothers 38% one or more children	Retrospective	18 weeks gestation	Energy and nutrient intake	Self-administered, validated FFQ (84 items)	Intake mean (no p-values reported) Energy (kcal) Pre-pregnancy 1852 (SD 751) Pregnancy 2104 (SD 583) Energy (MJ) Pre-pregnancy 7.75 (SD 3.14) Pregnancy 8.81 (SD 2.44) Carbohydrates (E%) Pre-pregnancy 48.1 (SD 5.3) Pregnancy 51.1 (SD 6.6) Protein (E%) Pre-pregnancy 14.6 (SD 2.1) Pregnancy 16.8 (SD 2.4) Fat (E%) Pre-pregnancy 35.9 (5.4) Pregnancy 32.1 (SD 6.4)	None Reported.
Crozier et al, 2009a (United Kingdom)	Sample size: 2057 Age: Not reported Gestation weeks: N/A Weight category: Not reported SES or similar: Not reported Education: Not reported Ethnicity: Not reported	Prospective	Pre-pregnancy, 11 and 34 weeks gestation. Conception was on average 1.8 years after pre-pregnancy data collection	White bread, breakfast cereals, cakes and biscuits, processed meat, crisps, fruit and fruit juices, dried fruit, sweet	Validated interviewer-administered FFQ	Intake of white bread, breakfast cereals, cakes and biscuits, processed meat, crisps, fruit and fruit juices, dried fruit, sweet spreads, confectionery, and hot chocolate drinks increased from pre-pregnancy to pregnancy (all p<0.0001). Consumption of breakfast cereals, cakes	None Reported. 24

	Smoking: Not reported NCD's: Not reported Parity: Not reported			spreads, confectionery, and hot chocolate drinks, fruit, sweet spreads, puddings, cream, milk, cheese, full-fat spread, cooking fats and salad oils, red meat, soft drinks, rice and pasta, liver and kidney, vegetables, vegetable dishes, nuts, tea, coffee, boiled potatoes, crackers.		and biscuits, processed meat, non-citrus fruit, sweet spreads, and hot chocolate drinks increased further in late pregnancy (all $p < 0.0001$). Puddings, cream, milk, cheese, full-fat spread, cooking fats and salad oils, red meat, and soft drinks did not change in early pregnancy, they increased in late pregnancy (all $p < 0.001$). Intakes of 10 foods or food groups decreased in pregnancy. These were consumption of rice and pasta, liver and kidney, salad vegetables, other vegetables, vegetable dishes, nuts, diet cola, tea, and coffee were lower in pregnancy than before pregnancy (all $p < 0.0001$). Compared to early pregnancy, consumption of rice, pasta, liver, and kidney were lower again in late pregnancy ($p < 0.001$). Consumption of green vegetables, boiled potatoes, and crackers did not change in early pregnancy but decreased in late pregnancy.	
Crozier et al, 2009b (United Kingdom)	Sample size: 1490 Age: 28.2 years Gestation weeks: N/A Weight category: Non-pregnant BMI mean 24.3 SES or similar:	Prospective	Pre-pregnancy, 11 and 34 weeks gestation	Portions of fruit and vegetables per day Caffeinated drinks/day (i.e. coffee,	Interviewer-administered 100-item food frequency questionnaire	Fruit and vegetable median scores 5.2 (IQR 3.7-7.0) pre-pregnancy, 5.3 (IQR 3.7-7.0) 11 weeks gestation, 5.4 (IQR 3.9-7.2) 34 weeks gestation. Eating <5 portions of fruit and	None Reported.

	Education: None 2.3% GCSE grade D or below 10.3% GCSE graded C or above 28.4% A level or equivalent 29.3% HND or equivalent 7.8% Degree 21.9% Ethnicity: 96.2% White 3.8% Non-white Smoking: 26.6% yes pre-pregnancy NCD's: Not reported Parity: Not reported			caffeinated tea and cola)		vegetables a day: 47% pre-pregnancy, 46% 11 weeks gestation, 44% 34 weeks gestation (NS change btw time points) Caffeinated drinks median scores: 4.1 (IQR 2-6) pre-pregnancy, 2.0 (IQR 0.6-4.1) 11 weeks gestation, 2.3 (IQR 0.9-4.3) 34 weeks gestation. Drinking >300mg of caffeine in drinks per day: 39% before pregnancy, 16% 11 weeks gestation, 20% 34 weeks gestation. All changes significant. P < 0.05	
Cuco et al, 2006a (Spain)	Sample size: 80 Age: 29 years (24-35 years) Gestation weeks: Not reported Weight category: 6.3% below BMI 20 70% BMI 20-25 20% BMI 25-30 3.8% above 30 BMI SES or similar: Not reported Education: Only primary education 22.5% Secondary education and vocational training 40% University education 37.5% Ethnicity: Not reported Smoking: Not reported NCD's: Not reported Parity: Not reported	Prospective	Pre-pregnancy, 6, 10, 26 and 38 weeks gestation and 6 months postpartum	Energy intake (kcal)	7 consecutive day dietary record	Data reported as 50 th percentile (25 th -75 th percentile) No p-values reported. Energy intake (kcal) Preconception 1910 (1730-2237) 6 weeks 1896 (1664-2076) 10 weeks 2017 (1743-2231) 26 weeks 2032 (1794-2251) 38 weeks 1899 (1680-2157) 6 months postpartum 1767 (1536-1957)	None Reported.
Cuco et al, 2006b(Spain)	Sample size: 77 Age: 27.3% 24-27 years 50.6% 28-31 years	Prospective	Pre-pregnancy, 6, 10, 26 and 38 weeks gestation	Energy intake (kcal), protein (g), carbohydrates	7 consecutive day dietary record	Data reported as 50 th percentile (25 th -75 th percentile) No p-values reported. Energy intake (kcal)	None Reported.

	22.1% ≥32 years Gestation weeks: N/A Weight category: pre-pregnancy: 6.5% <BMI 20 71.4% BMI 20-25 18.2% BMI >25-30 3.9% BMI >30 SES or similar: Not reported Education: Not reported Ethnicity: Not reported Smoking: 48.1% never 14.3% ex-smokers 13% pre-pregnancy NCD's: Not reported Parity: 67.5% primiparae			(g), fats (g), animal proteins (g) vegetable proteins (g)		Preconception 1940 (1743-2311) 6 weeks 1908 (1667-2084) 10 weeks 2037 (1742-2258) 26 weeks 2035 (1813-2299) 38 weeks 1904 (1688-2169) Proteins (g) Preconception 80 (71.5-91.9) 6 weeks 76.4 (68.8-86) 10 weeks 79.9 (67.3-87.6) 26 weeks 80.5 (70.6-93) 38 weeks 79.9 (68.4-87.5) Carbohydrates (g) Preconception 182.2 (157.3-226.4) 6 weeks 182.9 (163.1-212) 10 weeks 199.4 (178.9-230.2) 26 weeks 206.7 (175-239.9) 38 weeks 191.8 (165-223.4) Fats (g) Preconception 91.6 (82-118.2) 6 weeks 91.9 (79.8-103.6) 10 weeks 98 (79.8-110.1) 26 weeks 97.3 (83.8-111) 38 weeks 92.9 (75.6-104.6) Animal proteins (g) Preconception 54.7 (46.7-62.8) 6 weeks 51.7 (43.6-58.7) 10 weeks 48.5 (40.6-58.2) 26 weeks 50.9 (42.9-64.3) 38 weeks 52.9 (43.6-65.1) Vegetables proteins (g) Preconception 17.7 (14.6-22.9) 6 weeks 19.1 (15.5-22.7) 10 weeks 21.3 (16.6-25.5) 26 weeks 20.5 (17.4-24.9) 38 weeks 18.6 (15.9-22.2)	
Hellerstedt et	Sample size: 8827 (7174)	Retrospective	1-20 weeks	Daily Caffeine	Telephone	Caffeine:	Pregnancy intention:

al, 1997 (USA)	Age: 18-48yr Gestation weeks: Mean 8 weeks (1-20 weeks) Weight category: Not reported SES or similar: Employed - (82.7, 79.1, 68.8%) Education: 37% college degrees 12% graduate education Ethnicity: White – (89.1, 82.6, 77.7%) Smoking: Not reported NCD's: Not reported Parity: 65.5-94.9%		gestation		survey (yes/no, categorical questions)	<i>Preconception</i> (67.5, 69.8, 73.8) <i>Pregnancy</i> (26.0, 28.6, 38.7) All changes are $P < .01$.	Women with intended pregnancies, compared to those who reported the pregnancy was unintended, were more likely to report decreased consumption of caffeine in pregnancy.
Kopp-Hoolihan et al, 1999 (USA)	Sample size: 10 Age: 29.1 ± 5 (21-36 yrs) Gestation weeks: n/a Weight category: 23.1 ± 2.1 (19-26 kg/m ²) SES or similar: Not stated Education: Not stated Ethnicity: Not reported Smoking: Not reported NCD's: Not reported Parity: 2 nd or 3 rd child	Prospective	T0 - Preconception (within 3 months of pregnancy) T1, 2, 3 – Wk 8-10, 24-26, 34-36) TPost- 4-6 wk postpartum	RMR, DIT, TEE (active EE), EI and Body composition	3 day weighed food diary at each time point EI and Macronutrient content estimated at each time point from the 3d averaged values	Energy Intake only: 9% increase from T0 – T3 T0 – 8569 ± 1842 T1 – 8488 ± 1624 T2 – 8496 ± 1654 T3 – 9344 ± 2170 TPost – 8367 ± 2624 Large inter-individual variation	None reported

Paulik et al, 2009 (Hungary)	Sample size: 349 Age: 16-45 years Mean = 29.94 years Gestation weeks: 28.7 ± 0.7 weeks Weight category: not stated SES or similar: 7.4% Single Education: 37.5% secondary education 37.2% higher education Ethnicity: Not reported Smoking: Not reported NCD's: 78.7% in good or very good health Parity: 56.4% primiparae	Retrospective	During pregnancy (average 28.7 weeks gestation)	Fruit, vegetables, milk, coffee	Questionnaire	P-values not reported Fruit (85.7% vs 94.8%) Vegetables (67.6% vs 75.4%) Milk (66.8% vs 60.2%) Coffee (56.2% vs 33.2%)	None Reported
Pinto et al, 2008 Portugal	Sample size: 249 Age: 29 years (SD5.8) Gestation weeks: First trimester Weight category: 57% normal weight before pregnancy Pre-Preg BMI <18.5 = 3.4% 18.5-24 = 57.4% 25-30 = 28.4% >30 = 10.5% SES or similar: Employment- Student = 19.8%; employed = 59.5%; unemployed = 20.7%	Prospective	FFQ1 – first antenatal visit in trimester 1 (preconception) FFQ2 – After delivery (for whole pregnancy)	Energy (kcal) CHO (%TEI) Fat (%TEI) SFA (%TEI) Protein (%TEI) Caffeine (mg)	Semi-Quantitative FFQ with pre-specified portion sizes	Preconception vs pregnancy Energy (kcal) 2393 vs 2423 CHO (%TEI) 49.5% vs 50.3% Fat (%TEI) 31% vs 30.6% SFA (%TEI) 10% vs 10.5% Protein (%TEI) 17.6% vs 18.4% Caffeine (mg) 64.8 vs 34.4	None reported

	Education: <6yr = 31.7% 7-9yr = 29.3% 10-12yr = 26.1% >12 = 12.9% Ethnicity: Not stated Smoking: (1 st , 2 nd , 3 rd tri) (25%, 15.3%, 13.4%) NCD's: Not reported Parity: 0 = 62.7%; +1 = 37.3%						
Skreden et al, 2014 (Norway)	Sample size: 575 Age: 28.1 years (SD 4.35) Gestation weeks: 15 weeks gestation (range 5-20 weeks) Weight category: healthy weight (70.2%), overweight category (21.9%), obese category (7.5%) Mean BMI: 23.9 (SD 3.83) SES or similar: Not reported Education: 7-10 years 1.6% 10-12 years 12.9% Completed high school 16.9% < 4 years university/college 33.1% ≥ 4 yeayrs, 35.5% Ethnicity: Not reported Smoking: Not reported NCD's: Not reported Parity: Not reported	Retrospective	15 weeks gestation (range 5-20 weeks) And 'before they got pregnant'	Milk, water, coffee, sugar-sweetened beverages (SSB), artificially sweetened beverages (ASB), fruit juice	Food frequency questionnaire (0-never, 10-several times daily)	From pre-pregnancy to early pregnancy: the percentage of women drinking coffee decreased (38 % v. 10%, p<0.001), SSB decreased (10 % v. 6%, p=0.011) and ASB (12 % v. 9%, P =0.001) decreased of those reporting drinking it daily. Percentage of women who reported at least daily consumption of water (85 % v. 92%, P<0.001), fruit juice (14 % v. 20%, P=0.001) and milk (37 % v. 42%, P=0.001) increased.	Education: Women with higher educational attainment reduced their frequency of at least daily coffee consumption (46% v. 12%) more than women with lower educational attainment (31% v. 9 %; interaction timexeducation, P=0.005). Age (≥25 yrs vs <25yrs): Older women reported a larger decrease in at least daily consumption of artificially sweetened beverages (17% v. 11%) compared with younger women (7% v. 7 %; interaction timexage, (P=0.045).

							<p>Older women increased their frequency of at least daily consumption of fruit juice (17 % v. 27%) and daily intake of milk (35% v. 43%) from pre-pregnancy to early pregnancy more than younger women (fruit juice: 11% v. 13%; interaction time×age, $P=0.029$; milk; (39 % v. 40%; interaction time×age, $P=0.041$).</p> <p>BMI: No significant interactions found between BMI and changes in drinking habits from prepregnancy to pregnancy.</p>
Smedley et al, 2014 (Australia)	<p>Sample size: 100</p> <p>Age: 18 – 24 years = 11 >25 years = 89</p> <p>Gestation weeks: Postnatal (up to 12 months)</p> <p>Weight category: BMI: 18.5-25 = 69 25-30 = 20 30+ = 11</p> <p>SES or similar:</p>	Retrospective	12 months post birth (retrospective pre-conception and during pregnancy)	Dietary intake	Self-complete questionnaire (5-point Likert scale)	<p>Fruit (p 0.002) Never (10 v 5) Sometimes (25 v 17) Always (65 v 78)</p> <p>Veg (p 0.001) Never (10 v 5) Sometimes (29 v 18) Always (61 v 77)</p> <p>Fibre (p 0.001) Never (22 v 8) Sometimes (23 v 16)</p>	None reported

599	Employed = 91 Unemployed = 9 Education: High school = 20 Tech college = 22 University = 58 Ethnicity: n/a Country of Birth Oz = 70 Other = 30 Smoking: Non= 68 Ex = 18 Current = 4 NCD's: Not reported None Parity: 1 child = 88 1 + = 12					Always (55 v 76) Fried Food (NS) Never (56 v 67) Sometimes (34 v 28) Always (10 v 5) Fast Food (P0.017) Never (56 v 67) Sometimes (34 v 28) Always (10 v 5) Sweet Bakery (NS) Never (38 v 40) Sometimes (47 v 40) Always (10 v 20) Sweet Dairy (NS) Never (25 v 27) Sometimes (41 v 32) Always (34 v 41)	
-----	---	--	--	--	--	--	--

600 (SD=standard deviation, FFQ=food frequency questionnaire, NCD=Non-communicable diseases, RMR=Resting Metabolic Rate, DIT=Diet Induced Thermogenesis, TEE=Total
 601 Energy Expenditure, EI= Energy intake

602 **Supplementary Table 1:** Quality assessment of included studies

Author (year) / Checklist item	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	1.10	1.11	1.12	1.13	1.14	2.1
Aden et al, 2007	Yes	DNA	Yes	DNA	DNA	DNA	Yes	DNA	Can't Say	Yes	Yes	DNA	No	Yes	Acceptable
Crozier et al, 2009a	Yes	DNA	Yes	No	No	Yes	Yes	No	No	Yes	Yes	Yes	Can't Say	No	Acceptable
Crozier et al, 2009b	Yes	DNA	Yes	No	DNA	DNA	Yes	No	No	Yes	Yes	Yes	Can't Say	Yes	High quality
Cuco et al, 2006a	Yes	DNA	Yes	DNA	DNA	DNA	Yes	No	Can't Say	Yes	Can't Say	Yes	Yes	Yes	Acceptable
Cuco et al, 2005b	Yes	DNA	Yes	DNA	DNA	DNA	Yes	No	Can't Say	Yes	Can't Say	Yes	Yes	Yes	Acceptable
Hellerstedt et al, 1997	Yes	DNA	Yes	DNA	DNA	DNA	Yes	DNA	Can't Say	Yes	No	No	No	Yes	Acceptable
Kopp-Hoolihan et al, 1999	Yes	DNA	Yes	DNA	DNA	DNA	Yes	No	No	Yes	Can't say	Yes	Can't Say	No	Acceptable
Paulik et al, 2009	Yes	DNA	Yes	DNA	DNA	DNA	Can't Say	DNA	Can't Say	Can't Say	No	Yes	No	No	Acceptable
Pinto et al, 2009	Yes	DNA	Yes	DNA	DNA	DNA	Yes	DNA	Can't Say	Yes	Can't Say	DNA	No	Yes	Acceptable
Skreden et al, 2015	Yes	DNA	Yes	DNA	DNA	DNA	Yes	DNA	No	Yes	Can't say	DNA	Can't Say	Yes	Acceptable
Smedley et al, 2014	Yes	DNA	Yes	DNA	DNA	DNA	Yes	DNA	No	Can't Say	Can't Say	DNA	Can't Say	No	Acceptable

603 DNA – does not apply

604

605 **Checklist items**606 **1.1** The study addresses an appropriate and clearly focused question607 **1.2** The two groups being studied are selected from source populations that are comparable in all respects other than the factor under investigation.

608 (Deemed not applicable in this review)

609 **1.3** The study indicates how many of the people asked to take part did so, in each of the groups being studied610 **1.4** The likelihood that some eligible subjects might have the outcome at the time of enrolment is assessed and taken into account in the analysis.611 **1.5** What percentage of individuals or clusters recruited into each arm of the study dropped out before the study was completed? (Applies to prospective
612 studies only)613 **1.6** Comparison is made between full participants and those lost to follow up, by exposure status. (Applies to prospective studies only)614 **1.7** The outcomes are clearly defined.615 **1.8** The assessment of outcome is made blind to exposure status. If the study is retrospective this may not be applicable.616 **1.9** Where blinding was not possible, there is some recognition that knowledge of exposure status could have influenced the assessment of outcome617 **1.10** The method of assessment of exposure is reliable618 **1.11** Evidence from other sources is used to demonstrate that the method of outcome assessment is valid and reliable619 **1.12** Exposure level or prognostic factor is assessed more than once (In this review – has dietary intake been assessed more than once in
620 pregnancy/postpartum?)621 **1.13** The main potential confounders are identified and taken into account in the design and analysis.622 **1.14** Have confidence intervals been provided?623 **2.1** How well has the study done to minimise the risk of bias or confounding?

624

625 **Supplementary Table 2:** Details of studies excluded from systematic review and reason for exclusion.

Author	Year	Title and Journal	Reason for exclusion
Ådén et al.	2007	Energy and nutrients in self-reported diet before and at week 18-22 of pregnancy. <i>Scandinavian Journal of Food and Nutrition</i> 51(2): 67-73.	No diet data
Anderson et al.	2006	Prevalence of risk factors for adverse pregnancy outcomes during pregnancy and the preconception period -- United States, 2002-2004. <i>Maternal & Child Health Journal</i> 10(5): S101-106 101p.	No diet data
Arija et al.	2004	Food consumption, dietary habits and nutritional status of the population of Reus: Follow-up from preconception throughout pregnancy and after birth." <i>Medicina Clinica</i> 123(1): 5-11.	Manuscript not in English
Backhausen et al.	2014	Pregnancy planning and lifestyle prior to conception and during early pregnancy among Danish women. <i>European Journal of Contraception & Reproductive Health Care</i> 2014; 19 (1): 57-65.	No diet data
Bussell & Marlow	2000	The dietary beliefs and attitudes of women who have had a low-birthweight baby: a retrospective preconception study. <i>Journal of Human Nutrition & Dietetics</i> 13(1): 29-39 11p.	Between subject design
Clark & Ogden	1999	The impact of pregnancy on eating behaviour and aspects of weight concern. <i>International Journal of Obesity</i> . 23, 18±24	Between subject design
D'Angelo et al.	2007	Preconception and interconception health status of women who recently gave birth to a live-born infant -- Pregnancy Risk Assessment Monitoring System (PRAMS), United States, 26 Reporting Areas, 2004." <i>MMWR: Morbidity & Mortality Weekly Report</i> 56(SS-10): 1-35 35p.	No diet data
Grieger et al.	2016	"Asthma control in pregnancy is associated with pre-conception dietary patterns." <i>Public Health Nutrition</i> 19(2): 332-338 337p.	Between subject design
Harris et al.	2015	"Impact of rurality on maternal and infant health indicators and outcomes in Maine." <i>Rural & Remote Health</i> 15(3): 1-17 17p.	No diet data
Inskip et al.	2009	Women's compliance with nutrition and lifestyle recommendations before pregnancy: general population cohort study. <i>British Medical Journal</i> . 338:b481	Between subject design
Jedrychowski et al.	2007	Pre-pregnancy dietary vitamin A intake may alleviate the adverse birth outcomes associated with prenatal pollutant exposure: epidemiologic cohort study in Poland." <i>International Journal of Occupational & Environmental Health</i> 13(2): 175-180 176p.	No diet data
Kingsley et al	2012	Preconception health indicators among women - Texas, 2002-2010. <i>MMWR: Morbidity & Mortality Weekly Report</i> 61(29): 550-555 556p.	Between subject design

Oza-Frank et al.	2015	Provision of specific preconception care messages and associated maternal health behaviors before and during pregnancy." <i>American Journal of Obstetrics & Gynecology</i> 212(3): 372.e371-378 371p.	No diet data
Ramage et al.	2015	"Assessment of Pre-Pregnancy Dietary Intake with a Food Frequency Questionnaire in Alberta Women." <i>Nutrients</i> 7(8): 6155-6166 6112p.	Between subject design

626

627